

# Chronic Kidney Disease in the United States

---



U.S. Department of Health  
and Human Services



National Institutes of Health




National Institute of Diabetes and  
Digestive and Kidney Diseases

# Reasons for a National Kidney Disease Education Program

---


- 1) Kidney failure is a public health problem
- 2) Economical, effective testing and therapy exist
- 3) Testing and therapy are inadequately applied

# Kidney Failure is a Rapidly Growing Problem



# Kidney Failure Compared to Cancer Deaths in the U.S. in 2000

(in Thousands)




# Prevalence of Renal Insufficiency in U.S.

<b>GFR (mL/min/1.73 m<sup>2</sup>)</b>	<b>59-30</b>	<b>29-15</b>	<b>&lt; 15</b>
<b>Number of People</b>	<b>7.6 Million</b>	<b>360,000</b>	<b>&gt; 300,000</b>

**More than 8 million Americans have substantial kidney impairment and 10 million more have albuminuria.**


# Incident Rates by Primary Diagnosis

(per million population, adjusted)




# Incidence of Kidney Failure

(per million population, 1990, by HSA, unadjusted)



# Incidence of Kidney Failure

(per million population, 2000, by HSA, unadjusted)





# The Risk of Kidney Failure is Not Uniform

---

Relative risks compared to Whites:


*African Americans*     **3.9 X**

*Native Americans*     **2.9 X**

*Asians*     **1.6 X**

# **Costs of Kidney Failure are High**

**(in \$billions for 2000)**



# CVD is Linked to CKD

---

- Risk of CVD is increased 1.4 – 2.05 times with creatinine >1.4 – 1.5 mg/dl
- Risk of CVD is increased 1.5 – 3.5 times with microalbuminuria
- Annual mortality from CVD is increased 10 – 100 times with kidney failure
- First year CVD mortality (17%) is 5 times kidney failure incidence (3.5%) after diagnosis of CKD + diabetes

# Treatment to Prevent Progression of CKD to Kidney Failure

---

- Intensive glycemic control lessens progression from microalbuminuria in type 1 diabetes
  - DCCT, 1993
- Antihypertensive therapy with ACE Inhibitors lessens proteinuria and progression
  - Giatras, et al., 1997
  - Psait, et al., 2000
  - Jafar, et al., 2001

} **Meta-Analyses**
- Low protein diets lessen progression
  - Fouque, et al., 1992
  - Pedrini, et al., 1996
  - Kasiske, et al., 1998

} **Meta-Analyses**

# CKD is Not Being Recognized or Treated

---


- Most practices screen fewer than 10% of their Medicare patients with diabetes
- Patients are referred late to a nephrologist, especially African-American men
- Less than 1/3 of people with identified CKD get an ACE Inhibitor

# Is “System Level” Action Necessary?

---

- Universal medical coverage?
- Disease management teams?
- Improved reimbursement for prevention?
- Other?

# Age-Adjusted Cardiovascular Death is Declining



# Parallels Between Hypertension in 1972 and Kidney Disease in 2003

---

- Recent documentation of effective therapy
- Treatment of a silent disease to reduce risk for a disastrous outcome
- Simple screening
- Advantages for patients, physicians, industry



# Who to Test for Chronic Kidney Disease

---

Regular testing of people at risk

- Diabetes
- Hypertension
- Relative with kidney failure

# How to Test for Chronic Kidney Disease

---

- “Spot” urine albumin to creatinine ratio
- Estimate GFR from serum creatinine using the MDRD prediction equation

Note:

*24 hour urine collections are NOT needed*

*Diabetics: should be tested once a year*

*Others at risk: less frequently as long as normal*

# Who Should be Treated for Chronic Kidney Disease

---

- Diabetics with urine albumin/creatinine ratios more than 30mg albumin/1 gram creatinine
- Non-diabetics with urine albumin/creatinine ratios more than 300mg albumin/1 gram creatinine

**or**

- Non-diabetics with estimated GFR less than 60 mL/min/1.73 m<sup>2</sup>

# How to Treat for Chronic Kidney Disease

---

- Maintain blood pressure less than 130/80 mm Hg
- Use an ACE Inhibitor or ARB
- More than one drug is usually required and a diuretic should be part of the regimen


# How to Treat for Chronic Kidney Disease

(continued)

---

- Refer to dietician for a reduced protein diet
- Consult a nephrologist early
- Team with the nephrologist for care if GFR is less than 30 mL/min/1.73 m<sup>2</sup>
- Monitor hemoglobin and phosphorous with treatment as needed
- Treat cardiovascular risk, especially smoking and hypercholesterolemia

# Early Treatment Makes a Difference



# Target Audiences

---

- African Americans with
  - *Hypertension*
  - *Diabetes*
  - *Family history of kidney failure*
- Primary Care Providers

# NKDEP Activities

---

- “*You Have The Power To Prevent Kidney Disease*” awareness campaign
- Improved laboratory measurements and reporting of kidney function
- CKD quality indicators among Medicare beneficiaries hospitalized for cardiovascular disease
- Consult letter template for nephrologists
- Working with other non-profit, industry, and government groups



# PCP Must be Engaged

---

- 1) 7.6 million people with GFR 30-60 mL/min/1.73 m<sup>2</sup>
- 2) About 5,000 full-time nephrologists
- 3) Nearly 1,500 new patients per nephrologist

Therefore, 7 new patients per day per nephrologist.

Obviously not possible.

# What can Primary Care Providers do?

---

- Recognize who is at risk
- Provide testing and treatment
- Encourage labs to provide and report estimated GFR and spot urine albumin/creatinine ratios

# You Have The Power To Prevent Kidney Disease



# References

- Brenner BM, Cooper ME, de Zeeuw D, Keane WF, Mitch WE, Parving HH, Remuzzi G, Snapinn SM, Zhang Z, Shahinfar S, the RENAAL Study Investigators. Effects of Losartan on Renal and Cardiovascular Outcomes in Patients with Type 2 Diabetes and Nephropathy. *New England Journal of Medicine*. 2001 Sep 20;345(12):861-9.
- Collins AJ, Li S, Gilbertson DT, Liu J, Chen SC, Herzog, CA. Chronic Kidney Disease and Cardiovascular Disease in the Medicare Population. *Kidney International. Supplement*. 2003 Nov;(87):S24-31.
- Coresh J, Astor BC, Greene T, Eknoyan G, Levey AS. Prevalence of Renal Insufficiency in the U.S. *American Journal of Kidney Disease*. 2003 Jan;41(1):1-12.
- Flack JM, Neaton JD, Daniels B, Esunge P. Ethnicity and Renal Disease: Lessons from the Multiple Risk Factor Intervention Trial and the Treatment of Mild Hypertension Study. *American Journal of Kidney Disease*. 1993 Apr;21(4 Suppl 1):31-40.
- Gerstein HC, Mann JF, Yi Q, Zinman B, Dinneen SF, Hoogwerf B, Halle JP, Young J, Rashkow A, Joyce C, Nawaz S, Yusuf S; HOPE Study Investigators. Albuminuria and Risk of Cardiovascular Events, Death, and Heart Failure in Diabetic and Nondiabetic Individuals. *The Journal of the American Medical Association*. 2001 Jul 25;286(4):421-6.
- Jensen JS, Feldt-Rasmussen B, Strandgaard S, Schroll M, Borch-Johnsen K. Arterial Hypertension, Microalbuminuria, and Risk of Ischemic Heart Disease. *Hypertension*. 2000 Apr;35(4):898-903.

# References

## (continued)

- Kinchen KS, Sadler J, Fink N, Brookmeyer R, Klag MJ, Levey AS, Powe NR. The Timing of Specialist Evaluation in Chronic Kidney Disease and Mortality. *Annals of Internal Medicine*. 2002 Sep 17;137(6):479-86.
- Levey AS, Beto JA, Coronado BE, et al. Controlling the Epidemic of Cardiovascular Disease in Chronic Renal Disease: What do we Know? What do we Need to Learn? Where do we go from Here? National Kidney Foundation Task Force on Cardiovascular Disease. *American Journal of Kidney Disease*. 1998 Nov;32(5):853-906.
- McClellan WM, Ramirez SP, Jurkovitz C. Screening for Chronic Kidney Disease: Unresolved Issues. *Journal of the American Society of Nephrology*. 2003 Jul;14 (7 Suppl 2):S81-7. Review.
- McClellan WM, Knight DF, Karp H, Brown WW. Early Detection and Treatment of Renal Disease in Hospitalized Diabetic and Hypertensive Patients: Important Differences Between Practice and Published Guidelines. 1997 Mar;29(3):368-75.
- National Diabetes Information Clearing House. Diabetes Control and Complications Trial (DCCT). Bethesda (MD): National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, US Department of Health and Human Services; 1993 (NIH Publication No. 02-3874). Available from: <http://diabetes.niddk.nih.gov/dm/pubs/control/>
- Ries LAG, Eisner MP, Kosary CL, Hankey BF, Miller BA, Clegg L, Mariotto A, Fay MP, Feuer EJ, Edwards BK (eds). *SEER Cancer Statistics Review, 1975-2000*, National Cancer Institute. Bethesda, MD, [http://seer.cancer.gov/csr/1975\\_2000/](http://seer.cancer.gov/csr/1975_2000/),2003.

# References

## (continued)

---

- Ruilope LM, Salvetti A, Jamerison K, Hansson L, Warnold I, Wedel H, Zanchetti A. Renal Function and Intensive Lowering of Blood Pressure in Hypertensive Participants of the Hypertension Optimal Treatment (HOT) Study. *Journal of the American Society of Nephrology*. 2001 Feb;12(2):218-25.
- U.S. Renal Data System, USRDS 2003 Annual Data Report: Atlas of End-Stage Renal Disease in the United States, National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases, Bethesda, MD, 2003.
- U.S. Renal Data System, USRDS 2001 Annual Data Report: Atlas of End-Stage Renal Disease in the United States, National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases, Bethesda, MD, 2001.
- U.S. Renal Data System, USRDS 2000 Annual Data Report: Atlas of End-Stage Renal Disease in the United States, National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases, Bethesda, MD, 2000.